

CASE REPORT

Case Report: Moynihan's Hump: A Must Know Anatomical Variant for Every Laparoscopic Surgeon

Pranav Satyajit Wadhokar¹, Tejindersingh Tejpalsingh Chhabda², Satyajeet Gangadharrao Pathrikar³

Received on: 12 June 2022; Accepted on: 20 February 2023; Published on: 19 December 2023

ABSTRACT

Among the numerous anatomical variations in the hepatobiliary-vascular anatomy, Moynihan's hump, although rare, is a dangerous one. Failure of a surgeon to recognize it while operating can lead to complications like bile duct injury, bleeding and conversion to open surgery. We report a case of cholelithiasis, in which Moynihan's hump was identified during laparoscopic cholecystectomy and tackled with successfully. Further, we discuss the anatomy and surgical significance of this variation and conclude with the importance of adequate prior knowledge of hepato-biliary-vascular anatomy, achievement of critical view of safety and the use of indocyanine green dye as an adjunct to safely tackle this dangerous anomaly and prevent complications.

Keywords: Anomaly, Calot's triangle, Case report, Caterpillar hump, Laparoscopic cholecystectomy, Moynihan's hump, Right hepatic artery.

World Journal of Laparoscopic Surgery (2023): 10.5005/jp-journals-10033-1565

INTRODUCTION

Rigorous research has shown us the high percentage of variations in the anatomy of the hepato-biliary-vascular region, but even today, despite the use of modern techniques and imaging, the complications of iatrogenic injuries have not reduced.¹

Moynihan's hump is a rare (incidence of 3–13.3%) but disastrous variation, in which the right hepatic artery (RHA) has a tortuous course through the hepatocystic triangle, lying very close to the gallbladder with a short cystic artery. Misidentification of RHA as cystic artery can lead to inappropriate clipping, bleeding, or duct injury. In the literature, till date, the global proportion of Moynihan's hump in *in vivo* as well cadavers is only 7%.¹

CASE DESCRIPTION

A 44-year-old male presented with intermittent right hypochondriac pain of mild to moderate intensity and aggravating on eating, since the past 3 months with no comorbidities. An ultrasonogram of abdomen revealed cholelithiasis without any signs of cholecystitis. Patient underwent elective laparoscopic cholecystectomy for persistent biliary colic after preanesthetic fitness and consent. Patient was positioned in supine reverse Trendelenburg position. Pneumoperitoneum was created using Verees needle technique and subsequent trocars were inserted (anterior axillary, mid-clavicular, and epigastric) under vision. Initial visualization revealed a non-inflamed gallbladder with no adhesions. With adequate lateral traction at the Hartmann's pouch, Calot's triangle was dissected carefully at the level of infundibulum, just in line with sulcus of Rouviere. Medial and lateral peritoneum were dissected using monopolar hook to reveal a large tortuous artery running toward gallbladder and then away from it toward the liver. The size and the turning away of the artery from the gallbladder created a suspicion as to it being a tortuous RHA rather than cystic artery. On further gentle dissection using a monopolar hook, a small short cystic artery was seen arising from the tortuous RHA. Both the cystic artery and the RHA hump were anterior to the cystic duct (Fig. 1). Critical view of safety was

¹⁻³Department of General Surgery, MGM Medical College and Hospital, Aurangabad, Maharashtra, India

Corresponding Author: Pranav Satyajit Wadhokar, Department of General Surgery, MGM Medical College and Hospital, Aurangabad, Maharashtra, India, Phone: +91 7709135002, e-mail: pwadhokar94@gmail.com

How to cite this article: Wadhokar PS, Chhabda TT, Pathrikar SG. Case Report: Moynihan's Hump: A Must Know Anatomical Variant for Every Laparoscopic Surgeon. *World J Lap Surg* 2023;16(2):94–96.

Source of support: Nil

Conflict of interest: None

Patient consent statement: The author(s) have obtained written informed consent from the patient for publication of the case report details and related images.




Fig. 1: Intraoperative image showing the single loop U-shaped Moynihan's hump of RHA with a short cystic artery arising from it. Both the vessels lie anterior to the cystic duct

achieved demonstrating a tortuous RHA with a short cystic artery along with a cystic duct. The cystic artery was clipped carefully close to the gallbladder without excessive traction. Cystic duct was clipped and cut and retrograde cholecystectomy was carried out with adequate hemostasis. Postoperative stay was uneventful and patient was discharged on post-op day two. Patient was asymptomatic at 3 months of follow-up.

DISCUSSION

The anatomy of hepatobiliary triangle, that is, the biliary tree and cystic artery is complex due to three important reasons: (1) multiple vital structures (bile duct, hepatic artery, portal vein) lying in close approximation with each other, (2) multiple anatomical variations in biliary tree, cystic artery, and RHA, and (3) high incidence of these anatomical variations. These variations are as common as 20–50%.^{1–5} Knowledge about these variations is a must for surgeons before operating in this critical area as the course of cystic artery cannot be determined preoperatively and can only be established by careful dissection during operative procedure. Lack of knowledge is one of the major causes of avoidable complications like bile duct injury and vascular injury. Vascular injuries can result in the conversion of laparoscopic approach to open and the incidence is 0–1.9% with a mortality of 0.2%.^{3,6}

Moynihan's hump refers to the tortuous course of the RHA forming a U-shaped loop giving it a "caterpillar hump" appearance with an incidence of 3–13.3%.^{1,7,8} As far as the etiology is concerned, there have been multiple theories but none have been proven to be the exact cause. According to Taylor CR,⁹ architectural distortion due to cirrhosis of liver causes corkscrewing of intrahepatic arteries leading to a tortuous hepatic artery. As of now, no clinical evidence has been demonstrated to prove this hypothesis. According to Benson and Page,⁴ it is not an anomaly but artifact created due to traction during cholecystectomy. Similar incidence of this anomaly in operated individuals and in cadavers disproves this hypothesis. Miyaki¹⁰ suggested that the hump is embryological in origin. The embryonic liver is supplied by three segmental arteries arising from the aorta. The middle hepatic artery forms the future hepatic artery. The left and right fetal hepatic arteries become the accessory hepatic arteries from left gastric and superior mesenteric arteries in 25 and 18% of the cases. It is postulated that complete or partial persistence of the fetal arterial supply can lead to caterpillar hump.

Anatomically, the tortuous part of the RHA can either lie anterior to the common hepatic duct or posterior to it. The posterior location is more common (60%).¹¹ In our case, the artery lied anterior to hepatic duct. Depending on its tortuosity, the artery can course with a single loop (55% cases) when it is less tortuous and in double loop when it is more tortuous. The cystic artery is usually short whether it arises from single loop or double loop course, except when it arises from the proximal loop of the RHA.^{12,13} Such characteristics of this anatomical variation can easily lead to complications due to (1) misinterpretation of RHA as cystic artery, (2) mechanical/tractional injury to RHA/cystic artery during dissection. These factors can cause:

- Misinterpretation of RHA as cystic artery and its complete ligation can cause right lobar ischemic necrosis of liver.
- Misinterpretation of RHA as cystic artery with its partial ligation can cause hepatic artery pseudoaneurysm leading to uncontrolled bleeding.

- Vascular injury to RHA/cystic artery leading to bleeding which obscures the surgeon's vision and prompts blind coagulation leading to injury or complete transection of bile duct.
- Conversion to open cholecystectomy due to uncontrolled bleeding from vascular injury.

Such iatrogenic complications can be prevented by simple preventive measures which include:

- Acquisition of thorough knowledge of hepatobiliary anatomical variations by the surgeon.
- Achieving the critical view of safety before ligating any structures and ensuring only two structures are attached to gallbladder.
- When in doubt about the anatomy, or difficulty in dissection of calot's triangle, a second opinion should be taken from senior surgeon and priority to be given to patient safety and outcome.
- Use of IV bolus of indocyanine green dye to delineate cystic artery – hepatic artery anatomy during cholecystectomy.¹⁴

Clinical Significance

Moynihan's hump is a rare but dangerous anatomical variation of hepatic artery and can only be detected intraoperatively. Failure to identify it can lead to complications like uncontrolled bleeding, conversion to open surgery, and injury to bile duct. To avoid these iatrogenic complications, a surgeon must know these variations in hepatobiliary anatomy and follow the rule of critical view of safety strictly. It is necessary to understand that each cholecystectomy must be approached as a new case and the surgeon should be thoroughly vigilant keeping his mind and eyes open.

REFERENCES

1. Marano L, Bartoli A, Polom K, et al. The unwanted third wheel in the calot's triangle: Incidence and surgical significance of caterpillar hump of right hepatic artery with a systematic review of the literature. *J Min Access Surg* 2019;15:185–191. DOI: 10.4103/jmas.JMAS_75_18.
2. Tolino MJ, Tartaglione AS, Sturletti CD, et al. Variedades Anatómicas del Árbol Biliar: Implicancia Quirúrgica. *Int J Morphol* 2010;28(4): 1235–1240. DOI: 10.4067/S0717-95022010000400039.
3. Ding YM, Wang B, Wang WX, et al. New classification of the anatomic variations of cystic artery during laparoscopic cholecystectomy. *World J Gastroenterol* 2007;13(42):5629–5634. DOI: 10.3748/wjg.v13.i42.5629.
4. Benson EA, Page RE. A practical reappraisal of the anatomy of the extrahepatic bile ducts and arteries. *Br J Surg* 1976;63(11):853–860. DOI: 10.1002/bjs.1800631105.
5. Cimmino PT, Bocchetti T, Izzo L. Rilievi anatomo-chirurgici per la colecistectomia per via laparoscopica [Anatomo-surgical considerations in laparoscopic cholecystectomy]. *G Chir* 1992; 13(4):149–151.
6. Deziel DJ, Millikan KW, Economou SG, et al. Complications of laparoscopic cholecystectomy: A national survey of 4,292 hospitals and an analysis of 77,604 cases. *Am J Surg* 1993;165(1):9–14. DOI: 10.1016/s0002-9610(05)80397-6.
7. Kavitha KB. Dual cystic arteries in association with caterpillar hump of right hepatic artery – A case report and its surgical relevance. *J Clin Diagn Res* 2015;9(7):AD01–2. DOI: 10.7860/JCDR/2015/13089.6156.
8. Raghuvanshi DS, Ram M, Khan S, et al. Moynihan's hump: Our eyes see what our mind knows. *Indian J Surg* 2021;83:1077–1078. DOI: 10.1007/s12262-020-02535-x.
9. Taylor CR. Cirrhosis Imaging; 2015. Available from: <https://emedicine.medscape.com/article/366426-overview>.
10. Miyaki T. Patterns of arterial supply of the human fetal liver. The significance of the accessory hepatic artery. *Acta Anat (Basel)* 1989;136(2):107–111. DOI: 10.1159/000146806.

11. Dandekar U, Dandekar K, Chavan S. Right Hepatic Artery: A Cadaver Investigation and Its Clinical Significance. *Anat Res Int* 2015;2015:412595. DOI: 10.1155/2015/412595.
12. Jansirani D, Mugunthan N, Phalgunan V, et al. Caterpillar hump of right hepatic artery: Incidence and surgical significance. *Natl J Clin Anat* 2012;1:121124. DOI: 10.4103/2277-4025.298016.
13. Bhargava G, Singh H, Singh H, et al. Moynihan's hump of right hepatic artery: A case report. *CIBTech J Surg* 2014;3:42–44.
14. Boni L, David G, Mangano A, et al. Clinical applications of indocyanine green (ICG) enhanced fluorescence in laparoscopic surgery. *Surg Endosc* 2015;29(7):2046–2055. DOI: 10.1007/s00464-014-3895-x.